

Brane-localized Kinetic Terms in the Randall-Sundrum Model ^{*} [†]

H. Davoudiasl^{1,a}, J.L. Hewett^{2,b}, and T.G. Rizzo^{2,c}

¹*School of Natural Sciences, Institute for Advanced Study, Princeton, NJ 08540*

²*Stanford Linear Accelerator Center, Stanford, CA, 94309*

Abstract

We examine the effects of boundary kinetic terms in the Randall-Sundrum model with gauge fields in the bulk. We derive the resulting gauge Kaluza-Klein (KK) state wavefunctions and their corresponding masses, as well as the KK gauge field couplings to boundary fermions, and find that they are modified in the presence of the boundary terms. In particular, for natural choices of the parameters, these fermionic couplings can be substantially suppressed compared to those in the conventional Randall-Sundrum scenario. This results in a significant relaxation of the bound on the lightest gauge KK mass obtained from precision electroweak data; we demonstrate that this bound can be as low as a few hundred GeV. Due to the relationship between the lightest gauge KK state and the electroweak scale in this model, this weakened constraint allows for the electroweak scale to be near a TeV in this minimal extension of the Randall-Sundrum model with bulk gauge fields, as opposed to the conventional scenario.

^{*}Work supported in part by the Department of Energy, Contract DE-AC03-76SF00515

[†]e-mails: ^ahooman@ias.edu, ^bhewett@slac.stanford.edu, and ^crizzo@slac.stanford.edu

1 Introduction

The Randall-Sundrum (RS) model [1] offers a new approach to the hierarchy problem. Within this scenario, the disparity between the electroweak and Planck scales is generated by the curvature of a 5-dimensional (5-d) background geometry which is a slice of anti-de Sitter (AdS_5) spacetime. This slice is bounded by two 3-branes of equal and opposite tensions sitting at the fixed points of an S^1/Z_2 orbifold which are located at $\phi = 0, \pi$ where ϕ is the coordinate of the 5th dimension. The 5-d warped geometry then induces an effective 4-d scale Λ_π of order a TeV on the brane at $\phi = \pi$, the so-called TeV brane. Λ_π is exponentially smaller than the effective scale of gravity given by the reduced Planck scale, $\overline{M}_{Pl}$, with the suppression being determined by the product of the 5-d curvature parameter k and the separation r_c of the two branes, *i.e.*, $\Lambda_\pi = \overline{M}_{Pl} e^{-kr_c\pi}$. In this theory, the original parameters of the 5-d action are all naturally of the size $\sim \overline{M}_{Pl}$, while in the 4-d picture a hierarchy with $\Lambda_\pi/\overline{M}_{Pl} \lesssim 10^{-15}$ appears. It has been demonstrated [2] that this scenario is naturally stabilized without the introduction of fine-tuning for $kr_c \simeq 11$, which is the numerical value required to generate the hierarchy. This model leads to a distinct set of phenomenological signatures that may soon be revealed by experiments at the TeV scale; these have been examined in much detail [3, 4, 5].

For model building purposes, it is advantageous to extend the original framework of the RS model, where gravity alone propagates in the extra dimension, to the case where at least some of the Standard Model (SM) fields are present in the bulk [6]. The simplest such possibility is to have the SM gauge fields in the bulk [4, 5], while the other SM fields remain localized on the TeV brane. This scenario was examined in detail some time ago [4] where it was found that the couplings of the Kaluza-Klein (KK) excitations of the bulk gauge fields to matter on the boundary were enhanced by a factor of $\sqrt{2\pi kr_c}$ compared to those of the

corresponding zero-mode states. Since $kr_c \simeq 11$ as discussed above, this enhancement is numerically significant and is approximately a factor of $\simeq 8.4$. Severe constraints are then imposed on the gauge KK excitations from precision electroweak measurements; these imply that the mass of the lightest gauge KK state must be heavy with $m_1 \geq 25 - 30$ TeV. Due to the relation in this model between the masses of the KK states and the scale of physics on the TeV brane, this bound then correspondingly requires that $\Lambda_\pi \geq 100$ TeV. This places such a scenario in a very unfavorable light in terms of its original motivation of resolving the hierarchy.

Here, we investigate whether Brane Localized Kinetic Terms (BLKT's) for bulk gauge fields can modify these results. Recently, Carena, Tait and Wagner [7] examined the phenomenological influence of such localized kinetic terms for bulk gauge fields within the context of flat TeV^{-1} -sized extra dimensions. These authors showed that such terms can lead to significant modifications to the gauge KK spectrum, as well as to the KK state self-couplings and couplings to fields remaining on orbifold fixed points. Localized kinetic terms are expected to be present on rather general grounds in any orbifolded theory, as was shown by Georgi, Grant and Hailu [8], since they are needed to provide counterterms for divergences that are generated at one loop. In addition, these boundary terms can have important implications in a number of different situations such as model building [9], the construction of GUTs in higher dimensions with flat [10] or warped [11] geometries, and, in a different context, the quasilocalization of gravity [12]. Up to now the phenomenological implications of BLKT's in the case of warped extra dimensions have not been examined.

We examine the effects of localized kinetic terms in the case where the SM gauge fields are present in the RS bulk and the remaining SM fields are on the TeV-brane. As in the case of TeV^{-1} -sized extra dimensions, we find that these boundary terms alter both the KK spectrum as well as their associated couplings to the remaining wall fields, but in a manner

which is quite distinctive due to the curved geometry in this scenario. We will show that for a reasonable range of model parameters these modifications can be quite significant and lead to a natural reduction in the strength of the KK couplings. This allows the restrictions on Λ_π to be reduced by more than an order of magnitude and it would then no longer be unfavorable to have gauge fields be the only SM fields present in the RS bulk. The next section contains a detailed discussion of the modification of the usual gauge boson analysis within the RS framework due to the existence of brane localized kinetic terms. In Section 3 we discuss the implication of these results for the phenomenology of this scenario and our conclusions can be found in Section 4. Special limits of the KK mass equation are discussed in an Appendix.

2 Formalism

In this section, we derive the wavefunctions and the KK mass eigenvalue equation for bulk gauge fields with BLKT's in the RS scenario. We then compute the coupling of the gauge field KK modes to 4-d fermions localized on the TeV brane, where the scale of physics is of order the electroweak scale. Our derivation will demonstrate how the boundary terms modify these couplings from their original form and result in a possible softening of the precision electroweak bounds on the mass of the lightest KK gauge state. In the rest of the paper, KK modes refer to those from the bulk gauge fields, unless otherwise specified.

We perform our calculations for a $U(1)$ gauge field; the generalization to the case of non-Abelian fields is straightforward. Let us start with the 5-d action for the gauge sector

$$\begin{aligned}
S_A = & - \frac{1}{4} \int d^5x \sqrt{-G} \left\{ G^{AM} G^{BN} F_{AB} F_{MN} \right. \\
& \left. + [c_0 \delta(\phi) + c_\pi \delta(\phi - \pi)] G^{\alpha\mu} G^{\beta\nu} F_{\alpha\beta} F_{\mu\nu} \right\} ,
\end{aligned} \tag{1}$$

where the terms proportional to c_0 and c_π represent the kinetic terms for the branes located at $\phi = 0, \pi$, respectively, and the metric G_{MN} is given by the RS line element

$$ds^2 = e^{-2\sigma} \eta_{\mu\nu} dx^\mu dx^\nu - r_c^2 d\phi^2 \quad ; \quad \sigma = kr_c |\phi|. \quad (2)$$

Here the dimensionless constants $c_{0,\pi}$ would naturally be expected to be of order unity. Note that we can neglect terms proportional to $\partial_5 A_\mu$ on either brane as demonstrated by [7]. We will work in the $A_5 = 0$ gauge so that terms including A_5 can also be neglected. Thus only the usual 4-d kinetic term appears in the brane contributions to the action. In our notation, $A, B, \dots = 0, \dots, 3, 5$ and $\alpha, \beta, \dots = 0, \dots, 3$. Here, k is the 5-d curvature scale, r_c is the compactification radius, and $x^5 = r_c \phi$ with $-\pi \leq \phi \leq \pi$. The field strength is given by

$$F_{MN} = \partial_M A_N - \partial_N A_M. \quad (3)$$

The gauge field is assumed to have the KK expansion

$$A_\mu(x, \phi) = \sum_n A_\mu^{(n)}(x) \frac{\chi^{(n)}(\phi)}{\sqrt{r_c}}. \quad (4)$$

Substituting the above expansion in the action (1) yields

$$\begin{aligned} S_A = & - \frac{1}{4r_c} \int d^5x \sum_{m,n} \left\{ [1 + c_0 \delta(\phi) + c_\pi \delta(\phi - \pi)] F^{\mu\nu(m)} F_{\mu\nu}^{(n)} \chi^{(m)} \chi^{(n)} \right. \\ & \left. - (2/r_c^2) A^{\mu(m)} A_\mu^{(n)} e^{-2\sigma} [(d/d\phi) \chi^{(m)}] [(d/d\phi) \chi^{(n)}] \right\}, \end{aligned} \quad (5)$$

where now all the Lorentz indices are contracted using the 4-d Minkowski metric. In order to cast this action in the diagonal form, we demand

$$\int d\phi [1 + c_0 \delta(\phi) + c_\pi \delta(\phi - \pi)] \chi^{(m)} \chi^{(n)} = Z_n \delta^{mn} \quad (6)$$

and

$$\frac{1}{r_c^2} \int d\phi e^{-2\sigma} \left[\frac{d}{d\phi} \chi^{(m)} \right] \left[\frac{d}{d\phi} \chi^{(n)} \right] = Z_n m_n^2 \delta^{mn}. \quad (7)$$

These equations imply

$$\frac{d}{d\phi} \left(e^{-2\sigma} \frac{d}{d\phi} \chi^{(n)} \right) + [1 + c_0 \delta(\phi) + c_\pi \delta(\phi - \pi)] r_c^2 m_n^2 \chi^{(n)} = 0. \quad (8)$$

Away from the boundaries at $\phi = 0, \pi$, the solution to Eq.(8) is given by [4, 5]

$$\chi^{(n)}(\phi) = \frac{e^\sigma}{N_n} \zeta_1(z_n), \quad (9)$$

where $z_n(\phi) \equiv (m_n/k)e^\sigma$; $\zeta_q(z_n) \equiv [J_q(z_n) + \alpha_n Y_q(z_n)]$, with J_q and Y_q denoting the usual Bessel functions of order q . The normalization N_n is set by the condition

$$\int_{-\pi}^{+\pi} d\phi [\chi^{(n)}]^2 = 1, \quad (10)$$

implying that the zero mode is given by $\chi^{(0)} = 1/\sqrt{2\pi}$. Integrating Eq.(8) around the fixed points $\phi = 0$ and $\phi = \pm\pi$ yields

$$2 \frac{d}{d\phi} \chi^{(n)}(0) + c_0 r_c^2 m_n^2 \chi^{(n)}(0) = 0 \quad (11)$$

and

$$2 \frac{d}{d\phi} \chi^{(n)}(\pi) - e^{2kr_c\pi} c_\pi r_c^2 m_n^2 \chi^{(n)}(\pi) = 0, \quad (12)$$

respectively.

Let $\varepsilon_n \equiv z_n(0)$. For light KK modes with masses $m_n \sim 1$ TeV, which are of phenomenological interest, $\varepsilon_n \ll 1$; for $k \sim \overline{M}_{Pl}$ we have $\varepsilon_n \sim 10^{-15}$. Substituting for $\chi^{(n)}$ from

Eq.(9) in Eq.(11), and using the identity

$$x\zeta_0(x) = \zeta_1(x) + x\zeta_1'(x), \quad (13)$$

we obtain

$$\alpha_n = -\frac{J_0(\varepsilon_n) + \delta_0 \varepsilon_n J_1(\varepsilon_n)}{Y_0(\varepsilon_n) + \delta_0 \varepsilon_n Y_1(\varepsilon_n)}, \quad (14)$$

where $\delta_0 \equiv c_0 k r_c / 2$ and a prime (') denotes a derivative. Note that with c_0 of order unity, we may expect values of δ_0 of order 10 since kr_c takes on the value ~ 11 in order to address the hierarchy. Similarly, from Eq.(12), we obtain

$$\zeta_0(x_n) - \delta_\pi x_n \zeta_1(x_n) = 0, \quad (15)$$

where $\delta_\pi \equiv c_\pi k r_c / 2$ and $x_n \equiv z_n(\pi)$. The roots x_n of Eq.(15) yield the gauge KK mass spectrum, $m_n = x_n k e^{-kr_c \pi}$. We will study various limits of this root equation in the Appendix. Following the same arguments as above, we may expect that $|\delta_{0,\pi}| \lesssim \mathcal{O}(10)$, assuming no fine-tuning.

Due to the exponentially small value of ϵ_n , it may first appear that the δ_0 -dependent corrections to α_n are highly suppressed, and that α_n is small, of order ~ 0.01 , as in the standard RS scenario. However, an expansion of Eq. (14) in the parameter ε_n yields

$$\alpha_n \simeq -\frac{\pi/2}{\ln(x_n/2) - k r_c \pi + \gamma - \delta_0}, \quad (16)$$

where $\gamma \approx 0.577$ is Euler's constant. α_n remains small over most regions of parameter space of interest here, except for when $\delta_0 \simeq -kr_c \pi \simeq -35.4$. There are two sources of modifications to α_n as compared to the original RS framework due to the presence of the boundary terms: (1) the appearance of the term proportional to δ_0 in the denominator which is not suppressed by the exponential warp factor, and (2) the BLKT-modified values of x_n . In our numerical




Figure 1: (a) The behavior of the first five roots as δ_π is varied, assuming $\delta_0 = 0$. Note the sharp increase in x_n near $\delta_\pi = 0$ for all KK levels. (b) The large negative δ_π region is expanded to show the new root originating at the value $\delta_\pi = -kr_c\pi \simeq -35.4$.

analyses we will use the exact expression for α_n , rather than the approximate version given above, and we take $kr_c = 11.27$.

To get a flavor of how the KK tower mass spectrum is modified in this scenario with BLKT's, we must solve Eq.(15) for a range of values of $\delta_{0,\pi}$. We first take the simplest possibility by setting $\delta_0 = 0$, and examine the roots x_n as a function of δ_π . Our results are displayed in Fig. 1. From Fig. 1a, we see that in the region near $\delta_\pi = 0$ the roots undergo a substantial shift but remain relatively independent of variations in δ_π away from the origin. For large negative values of δ_π , the roots are almost the same as those obtained in the case of the graviton KK spectrum in the original RS framework [3]. This can be seen analytically by taking the limit of large $|\delta_\pi|$ in Eq.(15) and neglecting $\alpha_n \ll 1$. When the value of $-\delta_\pi$ exceeds $kr_c\pi$, a new root appears which is barely perceptible in Fig. 1a, but whose turn-on is shown in detail in Fig. 1b. The analytical source of this root is discussed in the Appendix. We will later see that this signals the appearance of an unphysical ghost, *i.e.*, a tachyonic state with negative norm.

When $\delta_0 \neq 0$ the modifications to the roots are sensitive to the sign of δ_0 . For $\delta_0 = \delta_\pi$, and in the range $\delta_\pi \gtrsim -30$, the roots appear to be very similar to those obtained when $\delta_0 = 0$, as is shown in Fig. 2a. The major difference in this case is the location of where the additional ghost root materializes; as shown in the Appendix, this root appears at the value $\delta_\pi = -(kr_c\pi + \delta_0)$. When $\delta_0 > (<)0$ the turn-on of this root thus moves to smaller (larger) values of δ_π . In the case $\delta_0 = \delta_\pi$, Fig. 2a shows that the new root appears at $\delta_\pi = -kr_c\pi/2$ as expected. We also see that for $\delta_\pi \lesssim -\pi kr_c$ the roots approach those which govern the KK graviton spectrum, as discussed above. In Fig. 2b, we display the roots for the case $\delta_0 = -\delta_\pi$. For most values of δ_π shown in the figure, these are found to be similar to the case where $\delta_0 = 0$ except that a new light root does not appear in this case. This is as expected, based on the discussion in the Appendix. When δ_π is large and positive,

$\delta_\pi \gtrsim \pi k r_c$, the roots again approach those for the case of KK gravitons, as anticipated.

When $\delta_\pi = 0$, the roots exhibit a weak dependence on δ_0 away from the $\delta_0 = -k r_c \pi$ region; this can be seen in Fig. 3. The δ_0 dependence of the roots is isolated in the coefficients α_n as seen in Eq.(16). From this we see that the roots have a slightly negative slope with increasing δ_0 .

3 Analysis

In this section, we study the couplings of the KK modes to the boundary fermions at $\phi = \pi$. The values of these couplings are the driving force behind the important constraints on the mass of the lightest KK gauge field in the original RS scenario [4]. In this section, we reexamine these bounds in the presence of brane localized kinetic terms. To begin, we note that the diagonalization conditions (6) and (7) yield the 4-d action

$$S_A = -\frac{1}{4} \int d^4x \sum_n Z_n \left(F^{\mu\nu(n)} F_{\mu\nu}^{(n)} - 2m_n^2 A^{\mu(n)} A_\mu^{(n)} \right), \quad (17)$$

where we have

$$Z_n = 1 + c_0 [\chi^{(n)}(0)]^2 + c_\pi [\chi^{(n)}(\pi)]^2. \quad (18)$$

Note that for physical fields, we demand that $Z_n > 0$ for all $n \geq 0$.

The coupling of a bulk gauge field A_μ to fermions ψ localized at $\phi = \pi$ is given by the action

$$S_\psi = - \int d\phi d^4x (det V) g_5 \bar{\psi} V_\alpha^\mu \gamma^\alpha A_\mu(x, \phi) \psi \delta(\phi - \pi), \quad (19)$$

where $V_\alpha^\mu = e^\sigma \eta_\alpha^\mu$ is the vielbein, $det V = e^{-4\sigma}$, and g_5 is the 5-d coupling constant. Using




Figure 2: (a) The behavior of the first five roots as a function of δ_π for the case $\delta_0 = \delta_\pi$. Here, the new root appears at the value $\delta_\pi = -kr_c\pi/2 \simeq -17.7$. (b) Same as above for the case $\delta_0 = -\delta_\pi$.




Figure 3: The behavior of the first five roots as a function of δ_0 taking $\delta_\pi = 0$.

the expansion (4) in Eq.(19), we obtain

$$S_\psi = - \int d^4x \left[\frac{g_5}{\sqrt{2\pi r_c}} \bar{\psi} \gamma^\mu A_\mu^{(0)} \psi + \sum_{n \neq 0} g_5 \frac{e^{\pi k r_c} \zeta_1(x_n)}{N_n \sqrt{r_c}} \bar{\psi} \gamma^\mu A_\mu^{(n)} \psi \right], \quad (20)$$

where we have performed the redefinition $\psi \rightarrow e^{3kr_c\pi/2}\psi$ to make the ψ kinetic terms canonical. Here, we have used the normalization

$$N_n^2 = \frac{x_n^2}{kr_c \epsilon_n^2} \zeta_1^2(x_n) W, \quad (21)$$

where

$$W = (1 - 2\delta_\pi + \delta_\pi^2 x_n^2) \left[1 - \frac{\epsilon_n^2 \zeta_1^2(\epsilon_n) [1 + 2\delta_0 + \epsilon_n^2 \delta_0^2]}{x_n^2 \zeta_1^2(x_n) [1 - 2\delta_\pi + x_n^2 \delta_\pi^2]} \right]. \quad (22)$$

Note that this normalization is mode dependent. To bring the gauge field kinetic terms in

Eq.(17) into the canonical form, we require

$$A_\mu^{(n)} \rightarrow \frac{A_\mu^{(n)}}{\sqrt{Z_n}}. \quad (23)$$

Eqs.(20) and (23) then yield

$$S_\psi = - \int d^4x \left[g_0 \bar{\psi} \gamma^\mu A_\mu^{(0)} \psi + \sum_{n \neq 0} g_n \bar{\psi} \gamma^\mu A_\mu^{(n)} \psi \right], \quad (24)$$

with

$$g_0 \equiv \frac{g_5}{\sqrt{2\pi r_c Z_0}} \quad (25)$$

and

$$g_n \equiv g_0 \sqrt{2\pi k r_c} \sqrt{\frac{Z_0}{W Z_n}}. \quad (26)$$

Here, g_0 is identified as the usual 4-d coupling constant for the interactions of the fermions with the zero-mode KK state. The KK excitation couplings are given by g_n , which due to the presence of the boundary terms are now mode dependent. Without the presence of the brane kinetic terms, one obtains $Z_0 = Z_n = 1$ which yields the result that g_n is an approximately mode-independent constant given by $g_n \simeq g_0 \sqrt{2\pi k r_c}$, since in this case $\alpha_n \sim 10^{-2}$ and $W \simeq 1$.

We can now compute the size of the effect of the brane localized kinetic terms on the ratio of couplings g_n/g_0 . Using Eq.(18), we obtain

$$Z_0 = 1 + \frac{\delta_0 + \delta_\pi}{\pi k r_c}, \quad (27)$$

and, neglecting terms of order ϵ_n ,

$$Z_n \simeq 1 + W^{-1} \left[2\delta_\pi + \frac{8\delta_0 \delta_\pi^2 \alpha_n^2}{\pi^2 \zeta_0^2(x_n)} \right]. \quad (28)$$




Figure 4: The ratios of the first five KK couplings to that of the zero-mode as a function of δ_π for the cases (a) $\delta_0 = 0$, (b) $\delta_0 = \delta_\pi$, and (c) $\delta_0 = -\delta_\pi$. The lightest (heaviest) KK mode corresponds to the top (bottom) curve in each case.

Note that similar to the case of TeV^{-1} -sized extra dimensions, Z_n is found to be n -dependent. From the requirement that $Z_0 > 0$, we obtain the constraint $\delta_0 + \delta_\pi > -\pi k r_c \simeq -35.4$. If this condition isn't satisfied, the state then has negative norm and is unphysical.

We first consider the case $\delta_\pi \geq 0$, in order to avoid the ghost states. Figures 4a-c display the existence of a significant fall-off of the ratio of couplings g_n/g_0 as a function of δ_π . The values of the coupling ratio in the negative δ_π region, away from the regions where the ghost states appear, are shown in Fig. 5a-c. In both cases we see the mode dependence of the coupling strength away from the value $\delta_\pi = 0$. Note that the coupling strength of the first KK excitation is substantially smaller when $\delta_\pi < 0$ compared to the case with positive values of δ_π . It is clear that compared to the original RS framework, the ratio g_n/g_0 is now naturally much reduced.

Due to the shifts in the mass spectrum and the reduced values of the ratio g_n/g_0 , we now expect that the inclusion of the localized kinetic terms can result in a significant loosening of the precision electroweak bounds [4] on the RS model. To obtain the constraints on the KK gauge boson mass spectrum, we need to consider the quantity

$$\tilde{V}(\delta_\pi, \delta_0) \equiv \sum_n \left(\frac{g_n}{g_0}\right)^2 \left(\frac{x_1}{x_n}\right)^2, \quad (29)$$

where the x_n are the roots given above, and examine its detailed dependence on $\delta_{0,\pi}$. This quantity describes the relative strength of the set of contact interactions induced by dimension-6 operators arising from KK gauge exchanges in precision electroweak observables [4]; a full description of the analysis procedure and the dependence of the precision electroweak variables on V can be found in Ref. [13]. Here, we perform a global fit to the most recent precision electroweak data sample [14] with the use of ZFITTER. Since for the case $\delta_{0,\pi} = 0$ we know from our previous analysis that precision measurements imply the bound $m_1|_{RS} \gtrsim 25 \text{ TeV}$ for the lightest gauge KK excitation [4], we can eas-




Figure 5: The ratios of the first five KK couplings to that of the zero-mode as a function of negative δ_π for the cases (a) $\delta_0 = 0$, (b) $\delta_0 = \delta_\pi$, and (c) $\delta_0 = -\delta_\pi$. The lightest (heaviest) KK mode corresponds to the top (bottom) curve in each case.

ily obtain the corresponding constraint for non-zero values of $\delta_{0,\pi}$. We obtain the bound $m_1|_{BLKT} \geq m_1|_{RS}[\tilde{V}(\delta_\pi, \delta_0)/\tilde{V}(0, 0)]^{1/2}$; the numerical results of this analysis is presented in Fig. 6a-b. Here, we see that the bound on m_1 falls rapidly as the magnitude of δ_π increases. For example, with $\delta_\pi \sim 10 - 20$, we see that m_1 can be as low as 2 – 3 TeV which will lead to an observable signal in Drell-Yan production for the first gauge KK excitation at the LHC. This is demonstrated in Fig 7 for the case of $m_1 = 5$ TeV. For larger values of δ_π , such states may be observable at the Tevatron. For $\delta_\pi < 0$ away from the ghost root region, the constraints from electroweak data are even weaker; here, direct searches at the Tevatron can place bounds on the parameter space.

So far, we have shown that the inclusion of boundary kinetic terms allow for a substantially lighter mass spectrum for the gauge KK states than in the conventional RS scenario. However, the question remains as to whether this ameliorates the stringent bound of $\Lambda_\pi \gtrsim 100$ GeV. To address this issue, we convert the constraint on m_1 into one on Λ_π by noting that $m_1 = x_1 \Lambda_\pi k / \overline{M}_{Pl}$, recalling that $k / \overline{M}_{Pl} \leq 0.1$, and using the values of x_1 that we computed above. Solving for Λ_π , we find the results displayed in Fig. 8. Note that for most values of the parameters, the resulting bound on Λ_π remains quite large for positive values of δ_π . This is due to the interplay of the modified roots and the constraints on the gauge KK mass spectrum. However, for $\delta_\pi < 0$, we see that the lower bound on Λ_π is easily reduced by almost an order of magnitude or more. This would seem to be the preferred region of parameter space for this model in its relation to the hierarchy problem.

4 Conclusions

A minimal, yet interesting, extension of the RS model is obtained by assuming that only gauge fields propagate in the bulk. However, this setup has been shown to be constrained by




Figure 6: (a) Lower bound on the mass of the lightest gauge boson KK excitation as a function of $\delta_\pi \geq 0$ from a fit to the precision electroweak data for the cases $\delta_0 = \delta_\pi$ (top/green), $\delta_0 = 0$ (middle/red), and $\delta_0 = -\delta_\pi$ (bottom/blue). (b) Same as above but now for the region $\delta_\pi \leq 0$ with $\delta_0 = \delta_\pi$ (bottom/green), $\delta_0 = 0$ (middle/red), and $\delta_0 = -\delta_\pi$ (top/blue).




Figure 7: Drell-Yan event rate at the LHC for the first gauge KK state in the RS scenario with boundary terms. We have assumed a luminosity of 300 fb^{-1} and have set $m_1 = 5 \text{ TeV}$. The steeply falling histogram represents the SM Drell-Yan continuum.

precision electroweak data to have a scale $\Lambda_\pi \gtrsim 100 \text{ TeV}$ for physics on the TeV brane [4]. This makes such a scenario less attractive as a means for resolving the gauge hierarchy. The stringent bound on the electroweak scale Λ_π is a result of the strong coupling of the gauge KK modes to the boundary fermions. It is thus interesting to examine if a modification of the boundary physics can lead to a relaxation of this constraint. One such possibility which is a simple extension of the theory, motivated by field theoretic considerations [8], is the addition of brane localized kinetic terms for the bulk gauge fields.

In this paper, we assumed the presence of localized gauge field kinetic terms on both the Planck and TeV branes in the RS model. We then derived the wavefunctions and the mass spectrum of the gauge KK modes. The couplings of the KK states to the SM brane localized fermions were then calculated. It was shown that for natural choices of parameters, a substantial suppression of these KK couplings, compared to the original model, can be




Figure 8: Bounds on Λ_π as a function of δ_π for the cases $\delta_0 = \delta_\pi$ (green), $\delta_0 = 0$ (red), and $\delta_0 = -\delta_\pi$ (blue), as derived from the constraints on the mass of the first gauge KK state. The color coding of the curves corresponds to $\delta_0 = \delta_\pi, 0, -\delta_\pi$ being represented by the top(bottom), middle(middle), bottom(top) curves on the right(left) handside of the figure.

achieved. We then reexamined the precision electroweak bounds on the lightest gauge KK mass or, equivalently, Λ_π . We found that for a reasonably wide range of parameters, the mass of the lightest gauge KK field can be as low as a few hundred GeV and hence will be visible at the LHC and possibly the Tevatron. The weakening of the constraints from precision data results in a significantly less severe bound on Λ_π ; one can now naturally obtain $\Lambda_\pi \lesssim 10$ TeV. This makes the RS framework with gauge fields in the bulk more favorable, in the context of a solution to the hierarchy problem, and facilitates the construction of more realistic models based on the RS proposal. In addition, for much of the $\delta_{0,\pi}$ parameter space, the gauge field and graviton KK excitations are nearly degenerate which may result in some interesting collider phenomenology.

Given the observations in this paper, we expect that the addition of brane localized kinetic terms for other bulk fields, such as gravitons, could significantly alter the phenomenology of the RS model, and could also lead to the appearance of new features in the low energy theory.

Acknowledgements

The work of H.D. was supported by the US Department of Energy under contract DE-FG02-90ER40542.

Note Added: A related article appeared at a similar time [15]; we thank these authors for pointing out a sign error in an earlier version of this manuscript.

Appendix

Here, we will show that near special values of the BLKT coefficients δ_0 and δ_π a vanishing root develops in Eq.(15), signaling the appearance of an unphysical ghost state. We will also discuss the large $|\delta_\pi|$ limit of this equation. In the limit where $x_n \rightarrow 0$, Eqs.(15)

and (16) yield

$$x_n^2 = \frac{-4(\delta_\pi + \delta_0 + kr_c\pi)}{-(kr_c\pi + \delta_0)(2\delta_\pi + 1) + (1 + \delta_\pi)}. \quad (30)$$

This equation is valid for $x_n \ll 1$, and for $\delta_\pi = -(\delta_0 + kr_c\pi)$, it yields $x_n^2 = 0$. Thus, for $\delta_\pi \leq -(\delta_0 + kr_c\pi)$, an unphysical ghost state, a state with negative norm, appears in the spectrum, as implied by Eq. (27). Note that in the original RS model, with $\delta_0 = \delta_\pi = 0$, this ghost state does not appear, since $kr_c\pi > 0$.

We now study Eq.(30) in the limit $\delta_\pi \rightarrow \pm\infty$, for the three cases $\delta_0 = 0, \delta_\pi, -\delta_\pi$ that were examined in the text. For $\delta_0 = 0$, we have $x_n^2 \rightarrow 4/(2kr_c\pi - 1)$ as $\delta_\pi \rightarrow \pm\infty$. Since $kr_c\pi \approx 35$, the small x_n approximation is valid and for $\delta_\pi \rightarrow +\infty$, the root is physical, as shown in Fig. 1a. Here, we see that the root for the lowest KK state asymptotes to a constant value of $\simeq 0.24$ as δ_π becomes large and positive; when δ_π is large and negative, the ghost root asymptotes to this same value. For $\delta_0 = \delta_\pi$, $x_n^2 \rightarrow 4/\delta_\pi$ as $\delta_\pi \rightarrow \pm\infty$. This root is only physical for $\delta_\pi \rightarrow +\infty$, and approaches zero, as presented in Fig. 2a. In the case where $\delta_0 = -\delta_\pi$, we have $x_n^2 \rightarrow -(4kr_c\pi)/(2\delta_\pi^2)$ as $\delta_\pi \rightarrow \pm\infty$, and there are no light physical modes in this limit, as seen in Fig. 2b.

We would like to note that in the limit $\delta_\pi \rightarrow \pm\infty$, with choices of δ_0 that yield $|\alpha_n| \ll 1$, Eq.(15) is well-approximated by $J_1(x_n) = 0$, for $x_n \neq 0$. In this case, apart from the special light modes discussed above, the rest of the gauge field KK modes have masses that are approximately degenerate with those of the KK gravitons from the original RS model [3]. More generally, as can be seen from Figures 1a and 2, a large range of parameters exist where this near degeneracy occurs. This will have important phenomenological implications at future colliders.

References

- [1] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221].
- [2] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999) [arXiv:hep-ph/9907447].
- [3] H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Rev. Lett. **84**, 2080 (2000) [arXiv:hep-ph/9909255].
- [4] H. Davoudiasl, J. L. Hewett and T. G. Rizzo, Phys. Rev. D **63**, 075004 (2001) [arXiv:hep-ph/0006041] and Phys. Lett. B **473**, 43 (2000) [arXiv:hep-ph/9911262].
- [5] A. Pomarol, Phys. Lett. B **486**, 153 (2000)[arXiv:hep-ph/9911294]; Y. Grossman and M. Neubert, Phys. Lett. B **474**, 361 (2000) [arXiv:hep-ph/9912408]; T. Gherghetta and A. Pomarol, Nucl. Phys. B **586**, 141 (2000); S. Chang, J. Hisano, H. Nakano, N. Okada and M. Yamaguchi, Phys. Rev. D **62**, 084025 (2000)[arXiv:hep-ph/9912498]; R. Kitano, Phys. Lett. B **481**, 39 (2000)[arXiv:hep-ph/0002279]; S. J. Huber and Q. Shafi, Phys. Lett. B **498**, 256 (2001) [arXiv:hep-ph/0010195] and Phys. Rev. D **63**, 045010 (2001) [arXiv:hep-ph/0005286]; S. J. Huber, C. A. Lee and Q. Shafi, arXiv:hep-ph/0111465; J. L. Hewett, F. J. Petriello and T. G. Rizzo, JHEP **0209**, 030 (2002) [arXiv:hep-ph/0203091]; F. Del Aguila and J. Santiago, arXiv:hep-ph/0111047, arXiv:hep-ph/0011143, Nucl. Phys. Proc. Suppl. **89**, 43 (2000)[arXiv:hep-ph/0011142] and Phys. Lett. B **493**, 175 (2000)[arXiv:hep-ph/0008143]; C. S. Kim, J. D. Kim and J. Song, arXiv:hep-ph/0204002; C. Csaki *et al.*, Phys. Rev. **D66**, 064021 (2002), hep-ph/0203034; G. Burdman, Phys. Rev. D **66**, 076003 (2002) [arXiv:hep-ph/0205329]; F. del Aguila and J. Santiago, arXiv:hep-ph/0212205.

- [6] W. D. Goldberger and M. B. Wise, Phys. Rev. D **60**, 107505 (1999) [arXiv:hep-ph/9907218].
- [7] M. Carena, T. Tait and C. Wagner, hep-ph/0207056.
- [8] H. Georgi, A.K. Grant and G. Hailu, Phys. Lett. **B506**, 207 (2001).
- [9] H.C. Cheng, K.T. Matchev and M. Schmaltz, hep-ph/0204342 and hep-ph/0205314.
- [10] L. Hall and Y. Nomura Phys. Rev. **D64**, 055003 (2001) and Phys. Rev. **D65**, 125012 (2002); Y. Nomura, Phys. Rev. **D65**, 085036 (2002); R. Contino, L. Pilo, R. Rattazzi and E. Trincherini, Nucl. Phys. **B622**, 227 (2002); M. Chaichian and A. Kobakhidze, hep-ph/0208129.
- [11] K. Agashe and A. Delgado, [arXiv:hep-th/0209212]; K. Agashe, A. Delgado and R. Sundrum, Nucl. Phys. B **643**, 172 (2002) [arXiv:hep-ph/0206099].
- [12] G.R. Davali, G. Gabadadze and M. Porrati, Phys. Lett. **B485**, 208 (2000); G.R. Davali and G. Gabadadze, Phys. Rev. **D63**, 065007 (2001); G.R. Davali, G. Gabadadze and M.A. Shifman, Phys. Lett. **B497**, 271 (2001); G.R. Davali, G. Gabadadze and M. Kolanovic, Phys. Rev. **D64**, 084004 (2001); S.L. Dubovsky and V.A. Rubakov, Int. J. Mod. Phys. **A16**, 4331 (2001); M. Chaichian and A.B. Kobakhidze, Phys. Rev. Lett. **87**, 171601 (2001); M. Carena *et al.*, Nucl. Phys. **609**, 499 (2001).
- [13] T. G. Rizzo and J. D. Wells, Phys. Rev. D **61**, 016007 (2000) [arXiv:hep-ph/9906234].
- [14] M. Grünewald, talk presented at the *31st International Conference on High Energy Physics*, Amsterdam, Netherlands, July 2002; LEP/SLD Electroweak Working Group, LEPEWWG/2002-01.

[15] M. Carena, E. Ponton, T. Tait and C. E. Wagner, arXiv:hep-ph/0212307.